

# COLLECTION AND ANALYSIS OF REMOTELY SENSED DATA FROM THE RHODE RIVER ESTUARY WATERSHED

## ANNUAL REPORT

October 10, 1971 to October 9, 1972

Prepared under Contract No. NAS6-1913 by  
CHESAPEAKE BAY CENTER FOR ENVIRONMENTAL STUDIES  
SMITHSONIAN INSTITUTION

For  
NATIONAL AERONAUTICS AND SPACE ADMINISTRATION  
WALLOPS STATION  
WALLOP ISLAND, VA. 23337

73  
(NASA-CR-62094) COLLECTION AND ANALYSIS  
OF REMOTELY SENSED DATA FROM THE RHODE  
RIVER ESTUARY WATERSHED Annual Report,  
10 Oct. 1971 - 10 Oct. 1972 (Smithsonian  
Institution) - 74 p HC \$6.75 CSCI 68H  
G3/13  
Unclass  
32002  
October

N74-18347


COLLECTION AND ANALYSIS OF REMOTELY SENSED DATA  
FROM THE RHODE RIVER ESTUARY WATERSHED


Annual Report

October 10, 1971 to October 9, 1972

Contract No. NAS6-1913  
National Aeronautics and Space Administration  
Wallops Station, Wallops Island, Virginia 23337

Chesapeake Bay Center for Environmental Studies  
Smithsonian Institution

  
Dale W. Jenkins, Ph.D.  
Co-Principal Investigator

  
Francis S. L. Williamson, Ph.D.  
Co-Principal Investigator

Staff

Daniel Higman, M.A., Botanist  
Susan Weck, M.A., Botanist

Consultant

David Challinor, Ph.D.

## TABLE OF CONTENTS

|   | Page |
|---|------|
| Introduction . . . . .  | 1    |
| Standardization of Autumn Coloration Changes with<br>Munsell Color Chips . . . . .            | 4    |
| Mapping Old Field Vegetation for Spray Irrigation<br>Project . . . . .                        | 16   |
| Interpretation and Verification of Salt Marsh Vegetation<br>Types by Remote Sensing . . . . . | 18   |
| Seasonality in the Marsh Vegetation of Rhode River . . . . .                                  | 22   |
| Analysis of Cover in Three Marsh Vegetation Types . . . . .                                   | 51   |
| Correlations of Cover, Spectral Signature, and Productivity<br>in Hog Island Marsh . . . . .  | 70   |
| References Cited . . . . .  | 72   |

**PRECEDING PAGE BLANK NOT FILMED**

## INTRODUCTION

The remote sensing study being conducted by the Smithsonian Institution and NASA, Wallops at the Smithsonian's Chesapeake Bay Center for Environmental Studies (the Center) underwent a major shift in emphasis during 1972, from the study of predominantly forest and field vegetation to that of salt marshes. The shift was made for three reasons: 1) in response to the greater emphasis placed by NASA on the problems of Chesapeake Bay; 2) to coincide with the need by the Chesapeake Research Consortium (CRC) for a wetland classification system; and 3) to promote the application of remote sensing by members of the Rhode River Program (RRP), a subgroup of the CRC working at the Center.

Before the marsh plants recommenced their annual growth in early summer, we concluded our work on the photointerpretation of forest and old field vegetation (begun in 1970-71). We sought a consistent basis for identifying species of forest trees by comparing natural color transparencies of their autumn foliage with standardized color chips from the Munsell Book of Color. Despite problems arising from small sample size and variations in illumination of the trees, the comparison with Munsell color chips proved to be a good identification technique for several species. This technique may also be feasible for separating and identifying species of salt marsh vegetation.

We also made a preliminary survey of the vegetation in an abandoned field and constructed a vegetation cover map. This survey anticipates that the site will be subjected to spray irrigation with secondarily treated sewage and that the consequent changes in the vegetation can be monitored by remote sensing.

Vegetation maps were constructed in early summer of all salt marshes in Rhode River (except those already mapped in 1971) by imposing current ground truth data on acetate tracings of available photographs. Since many marsh communities undergo seasonal variations in size, shape, and apparent composition, it was necessary to remap the marshes in the autumn. These seasonal changes made it difficult to predict the composition of all but the most homogeneous communities.

We spent most of the summer and autumn investigating and documenting the seasonality of salt marsh vegetation. In addition to repeated reconnaissance of all marshes in Rhode River, we closely examined the composition of three major plant communities in Hog Island marsh by stratified random sampling. From the samples we deduced the percentage of cover occupied by all species in these communities. We have not yet had time nor adequate equipment to compare the seasonal aspects of vegetation in each sample area. During 1972-73 we will broaden our sampling to include all marsh vegetation types.

We have attempted to determine how well our identification of marsh vegetation types by remote sensing will contribute to studies of primary productivity. Remote sensing has been used by Reimold (1971) to estimate the productivity of Georgia marshes. We invited Dr. Reimold to lecture at the Center about his methods and advise us as to their applicability to our marshes. We have used both Dr. Reimold's and Dr. Levin's (University of Pennsylvania) recommendations in correlating recent productivity measurements in Hog Island Marsh with our cover estimates and aerial signatures.

Our work in 1972-73 will concentrate on perfecting means of identifying marsh vegetation types by remote sensing, despite their seasonal variations. This work will be extended to marshes in other parts of the

Bay as necessary. After a reliable identification system has been developed, we will attempt to automate the recognition process by scanning microdensitometry. This method will be used to formulate a classification system for wetlands of the Chesapeake Bay.

## STANDARDIZATION OF AUTUMN COLORATION CHANGES WITH MUNSELL COLOR CHIPS

In order to correlate species differences with leaf coloration changes, observations of vernal and autumnal phenology were made on hardwood trees throughout the Rhode River watershed in 1971. Difficulties were encountered in correlating these observations with aerial photographs, because of vignetting and other color fidelity problems in the aerial films, and because of variation in color ranges between trees of the same species and within individual trees. Moreover, the relative rates of color change varied within a season and from year to year. To control these difficulties, it was decided to compare crown coloration patterns with standardized Munsell color chips, a technique described by Heller et al, (1964) and by Krumpke et al (1971). Krumpke's method of overcoming the problem of intraspecific color variation was to differentiate "cluster ranges" of the most common colors within the "phase" or more variable range of colors characteristic of a species. This method was chosen for use at the Chesapeake Bay Center.

Although both Heller and Krumpke used aerial photographs in their studies, it was decided to use 35mm ground photographs for at least the initial comparisons at the Chesapeake Bay Center, since these had proved to have better color fidelity and freedom from vignetting than the aerial films used. Moreover, it was essential to locate and identify precisely the individual trees being photographed. Since the entire crowns of trees had to be photographed, a 12 meter silo was chosen as the camera point. A series of overlapping photographs was taken around the horizon, using a Canon FT camera with Kodachrome II film (ASA 25), at approximately noon on clear days. One series was made during the third week in October and two more during the first and third weeks in November, to correspond with early, middle, and late stages of autumn coloration (autumn coloration was delayed in 1971 due to unusually warm weather).

The photographs were developed commercially as 35mm slides. A set of tracings was made from them (an example of which is given in Figure 1) and the trees identified by symbols (Table 1).

The Munsell book of color was used to make the comparisons. The Munsell book contains a large number of opaque color chips arranged by equally spaced divisions of hue, value, and chroma. (The hue of a color is its relation to red, yellow, green, blue, or purple; the value indicates its lightness, and the chroma its strength or departure from neutral). Under the Munsell system, a color is recorded by numbers assigned to these three attributes, e.g., "5Y 8/6" indicates that the color has a hue of 5 Yellow, a value of 8 (on a scale from 2 to 9), and a chroma of 6 (on a scale from 1 to 14).

Comparisons between the Munsell chips and the 35mm slides were made by exposing equal areas of a color chip and a slide to the same light source, with the light being transmitted through the slide and reflected off the chip. This is basically the same method used by Heller and Krumpe, but technical details differed since each study used different types of film and different light sources. Heller also made 35mm slides of the Munsell color chips so that both the standard colors and the colors being determined could be viewed by transmitted light, but this technique was omitted from the present study because of possible changes in color fidelity when the slides were developed.

Since neither Heller nor Krumpe described their comparative devices, it was necessary to design one for the present study. Figure 2 is a sketch of the instrument. It was built of plywood and aluminum, and measures 9.2 cm deep, 10 cm wide, and 12.1 cm high, not including the eye shield and base. The unit supports a 35mm slide and a Munsell color chip side by side about 5 cm above the base, so that light may shine through the lower window and be



FIGURE 1.





TABLE 1.

SYMBOLS USED FOR SPECIES OF TREES SHOWN IN FIGURE 1.

|    |  |
|----|--|
| Be | Box elder ( <i>Acer negundo</i> )            |
| C  | Cherry, black ( <i>Prunus serotina</i> )     |
| D  | Dogwood, flowering ( <i>Cornus florida</i> ) |
| J  | Juniper ( <i>Juniperus virginiana</i> )      |
| M  | Maple, red ( <i>Acer rubrum</i> )            |
| P  | Persimmon ( <i>Diospyros virginiana</i> )    |
| Q  | Oak ( <i>Quercus</i> sp.)                    |
| Qp | Pin oak ( <i>Quercus palustris</i> )         |
| Qs | Spanish oak ( <i>Quercus falcata</i> )       |
| S  | Sweetgum ( <i>Liquidambar styraciflua</i> )  |
| Sa | Sassafras ( <i>Sassafras albidum</i> )       |
| Sm | Sycamore ( <i>Platanus occidentalis</i> )    |
| T  | Tuliptree ( <i>Liriodendron tulipifera</i> ) |
| Wl | Willow, black ( <i>Salix nigra</i> )         |

Figure 2.

**COMPARATOR FOR MUNSELL CHIPS & 35MM. SLIDES**




reflected upward through the slide. The upper window permits light from the same source to reach the Munsell color chip. The light source, a 250-watt blue photoflood lamp, (No. B1 superflood BCA, GE or Sylvania), is available at most camera shops. The color temperature of this lamp is 4800<sup>0</sup> K, so that when supplemented by natural daylight it produces a true coloration on daylight-type color films. Since the Munsell chips are designed to be viewed under natural daylight, they should also render accurate color when viewed under the lamp with a natural daylight supplement. It was impractical to use direct sunlight for the light source, since optimal sunlight conditions are seldom available here in winter. With the lamp as the primary light source, variations in the background daylight, because of haze or cloud cover, are not considered significant.

To use the instrument, a 35mm slide is placed on the slide supports and all parts except the tree being viewed are masked with black paper. A color chip is selected with the aid of the Munsell selection of Charts for Plant Tissues. This chip is placed beside the slide and marked except for an opening about as large as the tree being viewed. The slide and chip are then viewed vertically by a single interpreter. While a stereoscope is impractical for this comparison, a large magnifying lens proves useful to accentuate the colored leaves from background elements in the picture.


The ranges of observed colors are shown in Figures 3 to 6. Only two of the three series of slides (for early and middle autumn colors) have been examined to date. The relatively small sample size prevented as detailed a distinction of hue and chroma values as Krumpke's study used, but in the case of sweetgum, black cherry, and tuliptree in the late October series, enough trees were included to distinguish "cluster ranges" in which leaf colors most frequently fell. A cluster range was also discernible for sweetgum in the

Figure 3.


# MUNSELL COLOR COMPARATIVE DIAGRAMS




Sweetgum  
Late October Series



Sweetgum  
Early November Series




Tuliptree  
Late October Series




Tuliptree  
Early November Series

Figure 4.


# MUNSELL COLOR COMPARATIVE DIAGRAMS




Black Cherry  
Late October Series



Black Cherry  
Early November Series




Red Maple  
Late October Series




Red Maple  
Early November Series

Figure 5.


# MUNSELL COLOR COMPARATIVE DIAGRAMS




Sassafras  
Late October Series



Sassafras  
Early November Series




Spanish Oak  
Late October Series




Spanish Oak  
Early November Series

Figure 6.



# MUNSELL COLOR COMPARATIVE DIAGRAMS



Virginia & Loblolly Pine  
Late October Series



Virginia & Loblolly Pine  
Early November Series





early November series. Although 19 species of trees are represented in the series, only 7 occurred frequently enough to permit meaningful color comparisons.

Since shaded portions of a crown proved to be about two hues darker and one step lower in value than unshaded portions, only trees which reflected direct sunlight on the photographs were selected for comparison with the Munsell chips. Variations in hue were even wider when part of a crown was back-lighted, but corresponding variations in value and chroma were relatively small. Several trees shown in both series exhibited enough side-lighting or back-lighting to eliminate them from evaluation, since they were located south of the camera point. Only slight variations in camera aperture were needed when the sequence of photographs was made, despite the movements of the camera in relation to the sun. Although Heller (1964) observed that chroma was susceptible to variation with differences in camera exposure, chroma variations within a species seldom exceed two steps. The exceptions were sweetgum and tuliptree, where variations were evenly distributed over four steps.

To measure color variation resulting from differences in lighting and camera exposure, color diagrams were drawn for loblolly and Virginia pine trees (Figure 6.). Although the actual coloration of these trees probably remained constant, pines in the late October series showed a variation of two steps in hue and three in value; in the early November series the variation was three steps in hue and three steps in value, with the hue moving toward a more yellow green color. This apparent variation in hue is difficult to explain, unless the proximity of hardwood trees with yellowing crowns contributed to a false evaluation. The variations in chroma may have resulted from camera exposure changes.

This exercise has shown that the use of Munsell color chips to define

the ranges of autumn leaf coloration in hardwood trees is a promising technique, but circumstances in the present study made it unreliable for definitive correlations between leaf color and species. A major reason for this lack of reliability is the uncertainty of color fidelity in photographic film processing; this often changes the apparent color of the same tree from one frame to the next. This problem might be overcome if standardized color chips or ground panels could be photographed on the same film as the vegetation, so that any loss of color fidelity in the film processing would be readily detectable.

A second major problem is reliable correlation between ground identifications of individual trees and aerial photographs of the forest canopy. In the present study, this problem was solved by using a ground-based camera point, but then there were additional problems of small sample size and extreme sensitivity to sun angle. Forest areas selected for interpretation must be well marked out on the ground and close coordination established between the airplane crew and ground observers.

A third problem, which hampers comparison of identification techniques and results by different investigators is the lack of a single standard light source and instrumentation. Consequently, the crown identification data collected by Heller (1964) and Krumpe (1971) may not be comparable with ours.

If these problems can be overcome, further experiments with Munsell color comparison should be undertaken using 22.9 x 22.9 cm transparencies instead of 35mm slides. A reliable means of identifying forest canopy coloration is greatly needed, and the Munsell technique is currently the only standard for comparison of colors.

## MAPPING OLD FIELD VEGETATION FOR SPRAY IRRIGATION PROJECT

In the spring, 1972, a decision was made to utilize abandoned fields at the CBCES for disposal of secondarily-treated domestic sewage by means of a spray irrigation system. Thus, there developed the opportunity to study the effects of controlled application of treated sewage on the vegetation of abandoned fields using remote sensing techniques. One prerequisite of such a study is the mapping and description of existing vegetation at the proposed site before any effluents are discharged. This preliminary work was completed in early June.

Almost all of the proposed irrigation site lies above the 15.24 m contour; it is relatively level, and has steep, forested sides sloping toward Muddy Creek and the Rhode River. Approximately half of its 1 hectare is proposed for irrigation.

The vegetation map (Fig. 7), drawn over a base map derived from aerial photographs, indicates the composition of the plant communities. Natural color photos taken in July (Flight 73) were best in showing the patterns of summer vegetation, and an infrared film taken in April (Flight 51) best showed the unpaved roads and pine stands masked during summer. The base map was prepared using a frame from the July film enlarged to a scale of approximately 1:3,850. An acetate tracing was then made of the vegetation patterns. A frame from the April film was similarly enlarged and superimposed on the tracing to show the roads.

Seven of the ten vegetation types in Fig. 7 represent variations of immature forest canopy, and three represent ground cover of unforested area. The vegetation types were described on the basis of those species visible on aerial photographs. A detailed analysis of the vegetation within each of the cover types, including biomass, density, and frequency, will be made when the irrigation system is built.

# PROPOSED SPRAY IRRIGATION SITE




Figure 7.

## INTERPRETATION AND VERIFICATION OF SALT MARSH VEGETATION TYPES BY REMOTE SENSING

The increased emphasis being given to the role of salt marshes, particularly in estuarine metabolism, by both NASA and investigators in the Rhode River Program, as well as the need to consolidate skills in the identification of marsh plants, acquired in 1971, prompted a new exercise in the interpretation and mapping of this vegetation. This has enabled us to evaluate the usefulness of photographs taken at different seasons for identifying species of marsh plants, and to check the accuracy of our own identifications. Vegetation maps of three salt marshes in Rhode River and one in West River, now in great demand by investigators in the Rhode River Program, were developed.

The four marshes chosen border Sellman, Bear Neck, and Fox Creeks in Rhode River, and Cheston Creek in West River. Although floristically similar, the composition and distribution of vegetation types is quite different. A similar study will be made of the remaining salt marshes in Rhode River later this summer.

The first step in the project was to trace all discernible vegetation patterns in each marsh from natural color photographs. Those taken in April (flight 51) and November (Rome flight 71-67) were selected to permit a comparison of seasonal differences in discernible patterns. The vegetation types and actual species composition of each were independently interpreted by Higman and Weck. Most of the predictions were made from the same frames of April and November films used for mapping, but the photos taken in June 1970 (flight 15) and July 1971 (flight 73) were also consulted.

The predicted vegetation types on the maps were then compared with the actual situation in the field, and detailed descriptions were made. The descriptions include estimated percentages of the areas covered by those species which should be visible on an aerial photograph. Criteria such as biomass, density, and frequency were not considered necessary; a more precise determination of the floristic composition can be made later. The field notes were then synthesized into descriptions of 20 vegetation cover types. Most of these were common to all marshes; thus the same pattern could be used to represent a particular type of vegetation on all maps.

While several contiguous parts of a marsh may have similar vegetation, as in the Sellman Creek and Cheston Point marshes, these parts differ in species composition to the extent that they are discernible on an aerial photograph. This difference does not justify designation as distinctive vegetation types.

The predictions made of the types of marsh vegetation were compared not only with the actual vegetation types, but with the films from which each prediction was originally based. Each investigator graded the other on the relative accuracy of each identification, and tables were made for each marsh. These results were combined (Table 2) to show the relative success of each observer in identifying 16 different vegetation types. We had little or no success in predicting four vegetation types not previously encountered. These are omitted from Table 2.

The data obtained from flight 51 (April) and Rome flight 71-67 (November) were most useful for the identification of vegetation types, since films from these had been selected for the base maps. Data from flights 15 (June) and 73 (July) were consulted only as supplements. The decision to use a particular film was made independently by each observer.

TABLE 2

## RELATIVE SUCCESS OF IDENTIFICATION OF SALT MARSH VEGETATION TYPES

-----  
 Observers: W=Weck, H=Higman  
 Numbers in parentheses are no. of correct or partial identifications  
 per no. of attempts using each film.

| Vegetation Type  | Flight 51 (Apr.)                     | Flight 15 (June)                 | Flight 73 (July)                     | Flight 71-67 (Nov.)                    |
|--|--------------------------------------|----------------------------------|--------------------------------------|--|
| <i>Phragmites communis</i><br>100%   | W: 100% (1 of 1)<br>H: not used      | H: 0% (0 of 2)                   | not used                             | W: 100% (1 of 1)<br>H: 0% (0 of 2)     |
| <i>Typha angustifolia</i><br>60-100%   | W: 0% (0 of 2)<br>H: 0% (0 of 4)     | W: not used<br>H: 0% (0 of 2)    | W: 33% (1 of 3)<br>H: 50% (2 of 4)   | W: 37% (3 of 8)<br>H: 20% (1.5 of 8)   |
| <i>Iva frutescens</i><br>80-100%   | W: 100% (1 of 1)<br>H: 100% (2 of 2) | W: not used<br>H: 66% (2 of 3)   | W: 100% (1 of 1)<br>H: 100% (1 of 1) | W: 80% (4 of 5)<br>H: 75% (3 of 4)     |
| <i>Iva</i> 60-90%, <i>Distichlis/Spartina patens</i> 10-40%                                | W: 75% (2 of 2)<br>H: 50% (1 of 1)   | W: not used<br>H: 90% (6.5 of 7) | W: 75% (2 of 2)<br>H: 75% (2 of 2)   | W: 75% (10 of 14)<br>H: 78% (14 of 18) |
| <i>Iva</i> 60-90%, <i>Distichlis/Scirpus olneyi</i> 10-40%                                 | W: 75% (1 of 1)<br>H: not used       | not used                         | not used                             | W: not used<br>H: 50% (2 of 2)         |
| <i>Iva</i> 20-50%, <i>Distichlis/Spartina patens</i> 50-80%                                | W: not used<br>H: 50% (1 of 2)       | W: not used<br>H: 50% (2 of 2)   | W: not used<br>H: 75% (2 of 2)       | W: 25% (2 of 4)<br>H: 50% (3 of 3)     |
| <i>Iva</i> 10-20%, <i>Spartina patens</i> 80-90%   | W: not used<br>H: 100% (1 of 1)      | not used                         | W: 0% (1 of 1)<br>H: 100% (1 of 1)   | W: 80% (1 of 1)<br>H: 25% (1 of 1)     |
| <i>Distichlis</i> and/or <i>Spartina patens</i> and/or <i>Eleocharis halophila</i> 70-100% | W: 50% (2 of 4)<br>H: 100% (2 of 2)  | not used                         | not used                             | W: 33% (2 of 6)<br>H: 30% (5 of 9)     |
| <i>Iva</i> 60-70%, <i>Spartina alterniflora</i> 10-20%                                     | W: not used<br>H: 65% (1 of 1)       | not used                         | W: 65% (1 of 1)<br>H: 65% (1 of 1)   | not used                               |
| <i>Spartina alterniflora</i><br>70-100%  | W: 30% (1 of 2)<br>H: not used       | W: not used<br>H: 75% (1 of 1)   | not used                             | W: 50% (1 of 2)<br>H: 50% (4 of 6)     |
| <i>Spartina alterniflora</i><br>50%, <i>S. patens</i> 50%                                  | not used                             | not used                         | not used                             | W: not used<br>H: 70% (1 of 1)         |
| <i>Scirpus olneyi</i> 60-90%,<br><i>Distichlis</i> 10-20%                                  | W: 40% (2 of 5)<br>H: 25% (1 of 1)   | W: not used<br>H: 25% (1 of 1)   | W: 0% (0 of 1)<br>H: not used        | W: 33% (2 of 6)<br>H: 20% (3 of 6)     |
| <i>Scirpus</i> 40-60%, <i>Distichlis/Spartina patens</i> 30-50%                            | W: not used<br>H: 50% (1 of 1)       | not used                         | W: 50% (1 of 1)<br>H: 30% (1 of 1)   | W: 75% (2 of 2)<br>H: not used         |
| <i>Iva</i> and/or <i>Baccharis halimifolia</i> 70-100%                                     | not used                             | W: not used<br>H: 50% (2 of 4)   | not used                             | W: 33% (1 of 3)<br>H: 20% (1 of 5)     |
| <i>Baccharis</i> 80-100%   | not used                             | not used                         | not used                             | W: not used<br>H: 0% (0 of 1)          |
| Trees & Shrubs   | W: 100% (1 of 1)<br>H: not used      | W: not used<br>H: 75% (3 of 4)   | not used                             | W: 50% (1 of 2)<br>H: 17% (1 of 6)     |

There seems to be relatively little difference in the value of films taken in April, July, and November for the identification of Iva frutescens and vegetation types in which it is prominent, judging by the success of both observers in identifying this species. The bushy habit of Iva is conspicuous. Low matted grasses (Distichlis spicata, Spartina patens) are best identified using the April film, although not enough identifications were made to make this conclusion certain. Photographs taken in April or November were equally useful for identifying Scirpus olneyi and Spartina alterniflora.

Difficulty was encountered in distinguishing short grass (Distichlis/ Spartina patens) from moderately tall grass (Spartina alterniflora) or sedge (Scirpus olneyi), especially since these species have been found in sympatry. Scirpus was occasionally confused with cattail (Typha angustifolia), or with dead Iva frutescens. In both instances, the photographs taken in November were more difficult to interpret.

Iva, Phragmites communis, and Typha angustifolia, in that order, were the most distinctive. Phragmites was identifiable on both the April and November films, while photos taken in July and November were best for Typha. Neither species occurred frequently or over large areas, except Typha in the Sellman Creek marsh. The communities of shrubs and stunted trees in the Bear Neck Creek and Cheston Point marshes were also distinctive, although sometimes difficult to distinguish from Iva frutescens.



## SEASONALITY IN THE MARSH VEGETATION OF RHODE RIVER

The marshes of Rhode River exhibit distinct vegetation changes within the summer growing season. When the marsh vegetation is fully leafed out, in early summer, the vegetation may either change color and texture or become obscured by late flowering perennials, annuals, or epiphytes. By late summer, once-familiar vegetation in the field and on aerial photos becomes less so to the observer. For this reason, it has become necessary to map the marshes at both ends of the growing season for accurate interpretation of aerial photographs and for adequate understanding of marsh vegetation dynamics. A generalized Rhode River marsh is presented in Figures 8 and 9. Figures 10 through 28 are maps of all of the major marshes at Rhode River, depicting seasonal variations observed.

Natural color transparencies were chosen from both early and late summer flights and the vegetation patterns for each marsh traced onto acetate for use as base maps in the field. These were then revised in the field. The primary changes in vegetation from early to late summer in Rhode River that can be detected on aerial photos are:

1. stem elongation and flowering of:

Typha angustifolia, Phragmites communis, Spartina cynosuroides

2. lush vertical growth, leafing, flowering, and reddening of stems of:

Acnida cannabina

3. thin to lush covering growth and flowering of:

Lythrum lineare, Ptilimnium capillaceum, Polygonum hydropiperoides,  
Mikania scandens, Cuscuta sp., Slirpus pbustus

4. spotty, sometimes clumped, vertical growth and flowering of:

Hibiscus palustris, Kosteletzkya virginica, Solidago sempervirens,  
Pluchea camphorata

5. lush, vertical growth, leafing, flowering: Panicum virgatum

THE MOST MARKED CHANGES IN THE INDIVIDUAL MARSHES IN RHODE RIVER ARE AS FOLLOWS:

|                        | <u>Early Summer</u>   | <u>Late Summer</u>   |
|------------------------|---|--|
| Hog Island Marsh:      | Small patch <u>Spartina alterniflora</u>  | Covered by <u>Ptilimnium cappillaceum</u> , <u>Lythrum lineare</u> , and <u>Polygonum hydropiperoides</u>      |
|                        | Large patch of <u>Spartina alterniflora</u>   | Covered by <u>Acnida cannabina</u>   |
|                        | Whole Marsh   | Interspersed with <u>Hibiscus palustris</u> , <u>Kosteletzkya virginica</u> , and <u>Solidago sempervirens</u> |
| Bear Neck Creek Marsh: | <u>Spartina alterniflora</u>  | Covered by <u>Hibiscus palustris</u> in flower   |
|                        | Whole marsh   | Interspersed with <u>Hibiscus palustris</u> , <u>Kosteletzkya virginica</u> , and <u>Solidago sempervirens</u> |
| Cheston Point Marsh:   | <u>Typha angustifolia</u> , <u>Iva frutescens</u> , and <u>Distichlis spicata/spartina patens</u> mixture | Overshadowed by fall flowering annuals and perennials  |
|                        | Whole marsh   | Interspersed with <u>Hibiscus palustris</u> , <u>Kosteletzkya virginica</u> and <u>Solidago sempervirens</u>   |
| Sellman Creek Marsh:   | Shrubs and dead trees   | Covered by <u>Mikania scandens</u>   |
|                        | <u>Scirpus Olneyi</u>   | Covered by <u>Scirpus robustus</u>   |
|                        | <u>Distichlis spicata/Spartina patens</u>   | Overgrown with <u>Scirpus Olneyi</u>   |
|                        | <u>Scirpus Olneyi</u>   | Covered by <u>Polygonum hydro-piperoides</u>   |
|                        | <u>Spartina alterniflora</u>  | Covered by <u>Acnida</u>   |
|                        | Whole marsh   | Interspersed with <u>Hibiscus palustris</u> , <u>Kosteletzkya virginica</u> , and <u>Solidago sempervirens</u> |

|                  |             |  |
|------------------|-------------|--|
| Fox Creek Marsh: | Whole Marsh | Interspersed with <u>Hibiscus palustris</u> , <u>Kosteletzkya virginica</u> , and <u>Solidago sempervirens</u> |
|------------------|-------------|--|

|                    |                             |
|--------------------|-----------------------------|
| Kirkpatrick Marsh: | No basis for comparison yet |
|--------------------|-----------------------------|

While the flowering Hibiscus palustris and Kosteletzkya virginica do not generally cover the vegetation around them in late summer, their large, conspicuous blossoms, especially in the former, can create puzzling textural as well as tonal changes in once-familiar vegetation patterns. Marsh goldenrod (Solidago sempervirens) can similarly confuse the photointerpreter unless he is aware of its presence; in small concentrated clumps, the flowers of this plant appear very bright yellow on natural color aerial transparencies.

It is easy to anticipate changes in color and texture due to the flowering of the ever-present tall marsh grasses, Typha, Phragmites, and Spartina cynosuroides. Plants such as Acnida cannabina, Pluchea camphorata, Lythrum lineare, Ptilimnium capillaceum, and Polygonum hydropiperoides, however, are cryptic until their late summer stem elongation and flowering. These plants appear tiny and insignificant in spring and early summer and are generally overlooked, indeed unseen, by the field observer estimating cover.

From our work here in Rhode River, we have found that knowledge of vegetation changes within the growing season is essential to identification of marsh types from aerial photographs and to understanding of marsh ecosystem dynamics.

**This page intentionally left blank**

FIGURE 8

## GENERALIZED RHODE RIVER MARSH












FIGURE 9  
GENERALIZED RHODE RIVER MARSH - KEY

|   |   | <u>Frequent Associates</u>   |
|---|---|--|
|    | <i>Typha angustifolia</i>                             | <i>Spartina cynosuroides</i> , <i>Hibiscus palustris</i>   |
|    | <i>Spartina alterniflora</i>                          | <i>Acnida cannabina</i> , <i>Spartina cynosuroides</i>   |
|    | <i>Distichlis spicata</i> /<br><i>Spartina patens</i> | <i>Scirpus Olneyi</i> , <i>Hibiscus palustris</i>  |
|   | <i>Scirpus Olneyi</i>                                 | <i>Distichlis spicata</i> / <i>Spartina patens</i>   |
|  | <i>Iva frutescens</i>                                 | <i>Distichlis spicata</i> , <i>Baccharis halimifolia</i> , <i>Spartina alterniflora</i>                      |
|  | <i>Phragmites communis</i>                            |  |
|  | <i>Panicum virgatum</i>                               | <i>Thelypteris palustris</i> , <i>Scirpus Olneyi</i> ,<br><i>Distichlis spicata</i> / <i>Spartina patens</i> |
|  | Shrubs and small trees                                | <i>Baccharis halimifolia</i> , <i>Panicum virgatum</i> ,<br><i>Thelypteris palustris</i>                     |

# HOG ISLAND MARSH - EARLY SUMMER ASPECT

















FIGURE 10

FIGURE 11  
HOG ISLAND MARSH - EARLY SUMMER KEY

-  *Typha angustifolia* 70 - 100%
  
-  *Spartina alterniflora* 60 - 100%
  
-  *Spartina alterniflora* 50%, *Iva frutescens* 20%, mud/water 20%,  
*Spartina cynosuroides* 10%
  
-  *Iva frutescens* 50 - 100%
  
-  *Spartina cynosuroides* 80 - 100%
  
-  *Iva frutescens* 40%, *Spartina cynosuroides* 40%
  
-  *Scirpus Olneyi* 70 - 90%
  
-  *Iva frutescens* 50%, *Scirpus Olneyi* 30%
  
-  *Distichlis spicata*/*Spartina patens* 60 - 100%
  
-  *Distichlis spicata*/*Spartina patens* 60%, *Scirpus Olneyi* 40%
  
-  *Phragmites communis* 80 - 100%
  
-  *Panicum virgatum* 80 - 100%
  
-  Shrubs and small trees



# DISTRIBUTION OF COVER SAMPLES

## HOG ISLAND MARSH - LATE SUMMER





















FIGURE 12

FIGURE 13

HOG ISLAND MARSH - LATE SUMMER KEY

-  *Phragmites communis* 80 - 100%
-  *Typha angustifolia* with scattered *Hibiscus palustris* 70 - 100%
-  *Distichlis spicata*/*Spartina patens* 60 - 100%
-  *Distichlis spicata*/*Spartina patens* 60%, *Scirpus Olneyi* 40%
-  *Scirpus Olneyi* 70 - 90%
-  *Acnida cannabina* 90 - 100%
-  *Spartina alterniflora* 60 - 100%
-  *Spartina alterniflora* 30 - 40%, *Iva frutescens* 10 - 20%, *Acnida cannabina* 20 - 40%
-  *Spartina cynosuroides* 80 - 100%
-  *Iva frutescens* 50 - 100%
-  *Iva frutescens* 40%, *Spartina cynosuroides* 40%
-  *Iva frutescens* 50%, *Scirpus Olneyi* 30%
-  *Panicum virgatum* 80 - 100%
-  *Ptilimnium capillaceum* 70 - 80%, *Polygonum hydropiperoides*/*Spartina alterniflora* 20 - 30%
-  *Polygonum hydropiperoides* 80%, *Spartina alterniflora* 20%
-  *Lythrum lineare* 60%, *Polygonum hydropiperoides* 40%
-  Shrubs and small trees

# KIRKPATRICK MARSH LATE SUMMER ASPECT




FIGURE 14

FIGURE 15  
KIRKPATRICK MARSH - LATE SUMMER KEY
















-  *Typha angustifolia* 80 - 100%
-  *Phragmites communis* 70 - 100%
-  *Panicum virgatum* 40 - 75%
-  *Spartina cynosuroides* 60 - 90%
-  *Spartina patens*/*Distichlis spicata* 60 - 100%
-  *Spartina patens*/*Distichlis spicata* 50%, *Panicum virgatum* 50%
-  *Scirpus Olneyi* 50 - 100%
-  *Scirpus Olneyi* 35 - 50%, *Spartina patens*/*Distichlis spicata* 35 - 50%
-  *Iva frutescens* 20 - 40%
-  *Iva frutescens* 40 - 75%
-  *Iva frutescens* 70%, *Distichlis spicata*/*Scirpus Olneyi* 30%
-  *Iva frutescens* 50%, *Distichlis spicata*/*Scirpus Olneyi*/  
*Phragmites communis* 50%
-  *Iva frutescens* 60%, *Spartina patens*/*Distichlis spicata* and  
*Spartina alterniflora* 40%
-  Shrubs and small trees

FIGURE 16

## BEAR NECK CREEK MARSHES



This page intentionally left blank

# BEAR NECK CREEK MARSH EARLY SUMMER ASPECT





FIGURE 17

FIGURE 18

BEAR NECK CREEK MARSH - EARLY SUMMER KEY





# BEAR NECK CREEK MARSH LATE SUMMER ASPECT

BEAR NECK CREEK

TREES

TREES



100 m

FIGURE 19

FIGURE 20

BEAR NECK CREEK MARSH - LATE SUMMER KEY



*Iva frutescens* 80 - 100%



*Iva frutescens* 40%, *Distichlis spicata*/*Spartina patens* 40%,  
*Solidago sempervirens* 20%



*Distichlis spicata*/*Spartina patens* 60 - 100%



*Distichlis spicata*/*Spartina patens* 60 - 70%, *Hibiscus palustris*  
20 - 30%, *Panicum virgatum* 10%



*Hibiscus palustris* 80 - 100%



*Scirpus Olneyi* 40 - 80%



*Distichlis spicata*/*Spartina patens* 60%, *Scirpus Olneyi* 40%



*Scirpus Olneyi* 40 - 50%, *Iva frutescens* 20 - 30%, *Distichlis*  
*spicata*/*Spartina patens* 20 - 40%



*Typha angustifolia* 90 - 100%



*Spartina alterniflora* 80 - 100%



*Phragmites communis* 100%



Shrubs and small trees



Mud/shallow water

This page intentionally left blank

FIGURE 21

## TRIBUTARY MARSHES OF BEAR NECK CREEK - LATE SUMMER ASPECT




FIGURE 22

# CHESTON POINT MARSH EARLY SUMMER ASPECT




FIGURE 23

CHESTON POINT MARSH - EARLY SUMMER KEY















-  *Typha angustifolia* 80-100%
-  *Typha angustifolia* 50%, *Distichlis spicata*/*Spartina patens* 50%
-  *Typha angustifolia* 60%, *Iva frutescens* 20%, *Distichlis spicata*/*Spartina patens* 20%
-  *Distichlis spicata*/*Spartina patens* 70-100%
-  *Distichlis spicata*/*Spartina patens* 60-80%, *Iva frutescens* 20-40%
-  *Iva frutescens* 60-90%, *Distichlis spicata*/*Spartina patens* 10-40%
-  *Panicum virgatum* 80-100%
-  *Iva frutescens*/*Baccharis halimifolia* 30%, *Panicum virgatum* 50%, *Hibiscus palustris*, *Solidago sempervirens*, and *Spartina patens* 20%
-  *Iva frutescens*/*Baccharis halimifolia* 70-100%
-  *Phragmites communis* 100%
-  *Spartina alterniflora* 80-100%
-  *Scirpus Olneyi* 100%
-  *Scirpus robustus* 50%, *Distichlis spicata*/*Spartina patens* 50%
-  Shrubs, small trees

FIGURE 24

# CHESTON POINT MARSH LATE SUMMER ASPECT


















FIGURE 25

CHESTON POINT MARSH - LATE SUMMER KEY

-  *Typha angustifolia* 80 - 100%
-  *Typha angustifolia* 50%, *Distichlis spicata*/*Spartina patens* 50%
-  *Distichlis spicata*/*Spartina patens* 70 - 100%
-  *Distichlis spicata*/*Spartina patens* 60 - 80%, *Iva frutescens* 20 - 40%
-  *Iva frutescens* 60 - 90%, *Distichlis spicata*/*Spartina patens* 10 - 40%
-  *Panicum virgatum* 80 - 100%
-  *Panicum virgatum* 50%, *Iva frutescens*/*Baccharis halimifolia* 30%, *Solidago sempervirens*, *Hibiscus palustris*, and *Spartina patens* 20%
-  *Iva frutescens*/ *Baccharis halimifolia* 70 - 100%
-  *Phragmites communis* 80 - 100%
-  *Spartina alterniflora* 80 - 100%
-  *Scirpus olneyi* 80 - 100%
-  *Scirpus robustus* 50%, *Distichlis spicata*/*Spartina patens* 50%
-  Shrubs, small trees
-  Mixed fall flowers, tall marsh grasses, mud



# SELLMAN CREEK MARSH EARLY SUMMER


100 m

TREES

TREES

SELLMAN CREEK

TREES



Mud/shallow water



*Spartina alterniflora* 50-80%



*Spartina alterniflora* 80-100%



*Distichlis spicata*/  
*Spartina patens* 60-100%



*Iva frutescens* 80-100%



*Iva frutescens* 20-40%  
*Distichlis spicata*/  
*Spartina patens* 60-80%



*Iva frutescens* 40-50%  
Shrubs and small trees 40-50%



*Scirpus Olneyi* 50-80%



*Typha angustifolia* 60-100%



*Hibiscus palustris* 50%  
*Typha angustifolia* 50%



Fresh water marsh: grasses,  
mints, arrowhead, shrubs




Shrubs and dead trees

# **SELLMAN CREEK MARSH LATE SUMMER**

100 m

SELLMAN CREEK



TREES

TREES

TREES



FIGURE 27

FIGURE 28

# FOX CREEK MARSH - ALL SUMMER



*Scirpus Olneyi* 70-80%, *Distichlis spicata* 20-30%



*Scirpus Olneyi* 50%, *Distichlis spicata* 50%



*Iva frutescens* 60-70%, *Spartina alterniflora* 30-40%



*Typha angustifolia* 100%



*Iva frutescens* 60-90%, *Spartina patens*/  
*Distichlis spicata* 10-40%




*Iva frutescens* 90-100%, *Spartina patens* 0-10%



*Spartina patens* 80-90%, *Iva frutescens* 10-20%



*Spartina cynosuroides* 100%



## SEASONAL VARIATION IN PHRAGMITES COMMUNIS - A SPECIAL CASE

Patches of Phragmites communis are easily seen both on the ground and in aerial photographs; the plant grows in pure stands and retains the flowering head on dead stems through winter and spring until new growth replaces it in mid to late summer. On natural color aerial transparencies, Phragmites patches look the same; gray or blue-gray with a smooth, thick texture. On aerial photos taken in late summer, however, the largest patch of Phragmites in Rhode River, that on Kirkpatrick Marsh, appears brown. The small patches of Phragmites on Hog Island, Bear Neck Creek, and even Kirkpatrick Marsh appear typically gray or blue-gray. Flights 32 (11/8/70) and 89 (10/7/71) both show this difference. The appearance of Phragmites on different films is given below.

| <u>Flight</u> | <u>Date</u> | <u>Film Type</u> | <u>Large Patch<br/>in Kirkpatrick</u> | <u>Small Patches</u> |
|---------------|-------------|------------------|---------------------------------------|----------------------|
| 15            | 6/30/70     | Nat. Color       | Blue-gray                             | Blue-gray            |
| 32            | 11/8/70     | Nat. Color       | Light brown                           | Gray                 |
| 73            | 7/13/71     | Nat. Color       | Blue-gray                             | Blue-gray            |
| 89            | 10/7/71     | Nat. Color       | Dark brown                            | Gray                 |
| 113           | 6/7/72      | Nat. Color       | Blue-green-gray                       | Blue-gray            |

The individual Phragmites plants making up the large patch are markedly less robust than those in the smaller patches. The later are 2.5 to 3 meters tall with flowering heads of 33 x 15 cm while the plants in the large patch are 1.5 to 1.8 meters tall with flowering heads of 13 x 5 cm. It would seem that such a striking difference in vigor would result in color differences on aerial photos throughout the year. At this time, the cause of the lessened vigor in the large Phragmites patch in Kirkpatrick marsh is unknown.

The fact that such changes can occur seasonally with vegetation types thought to be readily identifiable indicates that seasonal monitoring is essential for accurate photointerpretation.

## ANALYSIS OF COVER IN THREE MARSH VEGETATION TYPES

The seasonal variability of marsh vegetation has necessitated the accurate determination of marsh composition and cover patterns. To date, this determination has been made for three major vegetation types or communities in Hog Island Marsh: those dominated by Iva frutescens, Distichlis spicata/Spartina patens, and Typha angustifolia. These communities were chosen for their size and ease of recognition on natural color aerial transparencies. Their boundaries had previously been mapped. Hog Island Marsh was chosen as a site because of its accessibility, the presence of a 100 m grid system and its recent use for productivity studies.

The three communities were divided into sections of approximately 20 m<sup>2</sup> each. Three sampling sites were chosen in each section, using a random numbers table. This method produced a fairly even "stratified" coverage of each community while still allowing statistical testing of the data. The sampling sites were marked on a 10 m<sup>2</sup> grid superimposed over an aerial transparency (Fig.12). Percentage of cover for each plant species and for exposed mud or water was estimated at each site with a 1 X 2 m sampling frame. The cover of the deciduous Iva frutescens was estimated as if all the leaves were present.

Figures 29 - 42 show the composition and percentages of cover in each sample plot. Table 3 shows the mean percentage of cover, standard deviation, and standard error of the mean for all species in each community.

As indicated by the standard deviation in Table 3, the Typha community is least heterogeneous with respect to cover variation within the community. The Distichlis/Spartina community is intermediate, with the Iva community most heterogeneous. The denseness of the Distichlis/Spartina community is reflected in the high percentage of area covered by the type species association (70 percent) while exposed mud and water accounted for only 8 percent.

The Iva frutescens community is by far the most heterogeneous as to species composition, although many of the associated species account for little cover. Cover variation within the community is also very high; this is evidenced by the high standard deviation in species with most cover: Iva frutescens, Distichlis spicata/Spartina patens, and Scirpus Olneyi.

As shown in Figures 29 - 42, the eastern and western stands of the Iva community in Hog Island marsh differ markedly in the percent cover of the type species: the eastern stand has a mean Iva cover of 33 percent while the western stand has a mean Iva cover of 16 percent. The latter case is interesting in that it can still be recognized as an Iva community with such a low percent cover of the type species. The highest percent cover is contributed by mud and shallow water. The significant area of marsh covered by mud and water, especially in the Iva community, demonstrates that texture and color seen on aerial photographs may be due as much to the spaces between plants as to the plants themselves.

TABLE 3

MEANS, STANDARD DEVIATIONS, AND STANDARD ERRORS OF COVER SAMPLES

| <u>Vegetation Type</u>                 | <u>n Sample Number</u> | <u>Species Present</u>                 | <u>Mean Cover (% 2-m<sup>2</sup> plot)</u> | <u>S.D. Standard Deviation</u> | <u>S.E. Standard Error of the Mean</u> |
|--|------------------------|--|--|--------------------------------|--|
| Typha                                  | 10                     | Typha angustifolia                     | 56   | 8                              | 3                                      |
|  |                        | Polygonum hydropiperoides              | 2  | 3                              | 1                                      |
|  |                        | Spartina alterniflora                  | 1  | 2                              | 0.7                                    |
|  |                        | Hibiscus palustris                     | 1  | 2                              | 0.7                                    |
|  |                        | Iva frutescens                         | 2  | 6                              | 2                                      |
|  |                        | Scirpus Olneyi                         | 5  | 10                             | 3                                      |
|  |                        | Mud/water                              | 29   | 11                             | 4                                      |
| Distichlis spicata/<br>Spartina patens | 15                     | Distichlis spicata/<br>Spartina patens | 70   | 21                             | 5                                      |
|  |                        | Scirpus Olneyi                         | 17   | 14                             | 4                                      |
|  |                        | Acnida cannabina                       | 0.3  | 0.7                            | 0.2                                    |
|  |                        | Solidago sempervirens                  | 1  | 1.5                            | 0.4                                    |
|  |                        | Polygonum hydropiperoides              | 1  | 1                              | 0.4                                    |
|  |                        | Mud/water                              | 8  | 10                             | 2.5                                    |
|  |                        |  |  |                                |  |
| Iva frutescens 27                      | 27                     | Iva frutescens                         | 25   | 18                             | 3                                      |
|  |                        | Distichlis spicata/<br>Spartina patens | 15   | 20                             | 4                                      |
|  |                        | Scirpus Olneyi                         | 14   | 14                             | 3                                      |
|  |                        | Polygonum hydropiperoides              | 2  | 5                              | 0.9                                    |
|  |                        | Spartina cynosuroides                  | 8  | 13                             | 3                                      |
|  |                        | Spartina alterniflora                  | 4  | 7                              | 1                                      |
|  |                        | Solidago sempervirens                  | 0.07                                       | 0.4                            | 0.1                                    |
|  |                        | Hibiscus palustris                     | 0.6  | 2                              | 0.4                                    |



TABLE 3 (Continued)

| <u>Vegetation<br/>Type</u>        | <u>n<br/>Sample<br/>Number</u> | <u>Species<br/>Present</u>             | <u>Mean Cover<br/>(% 2-m<sup>2</sup> plot)</u> | <u>S.D.<br/>Standard<br/>Deviation</u> | <u>S.E. Standard<br/>Error of<br/>the Mean</u> |
|-----------------------------------|--------------------------------|--|--|--|--|
| Iva frutescens                    | 27                             | Pluchea camphorata                     | 1  | 4                                      | 0.7  |
|                                   |                                | Kosteletzkya virginica                 | 0.6  | 2                                      | 0.3  |
|                                   |                                | Baccharis halimifolia                  | 0.4  | 2                                      | 0.4  |
|                                   |                                | Juncus spp.                            | 0.4  | 1                                      | 0.3  |
|                                   |                                | Panicum virgatum                       | 4  | 9                                      | 2  |
|                                   |                                | Mud/water                              | 25   | 17                                     | 3  |
| Iva frutescens<br>(Eastern plots) | 15                             | Iva frutescens                         | 33   | 20                                     | 5  |
|                                   |                                | Distichlis spicata/<br>Spartina patens | 20   | 22                                     | 6  |
|                                   |                                | Scirpus Olneyi                         | 8  | 11                                     | 3  |
|                                   |                                | Polygonum hydropiperoides              | 2  | 4                                      | 1  |
|                                   |                                | Spartina cynosuroides                  | 5  | 11                                     | 3  |
|                                   |                                | Spartina alterniflora                  | 6  | 9                                      | 2  |
|                                   |                                | Hibiscus palustris                     | 0.3  | 1                                      | 0.3  |
|                                   |                                | Pluchea camphorata                     | 0.9  | 3                                      | 0.7  |
|                                   |                                | Panicum virgatum                       | 6  | 11                                     | 3  |
|                                   |                                | Mud/water                              | 24   | 19                                     | 5  |
| Iva frutescens<br>(Western plots) | 12                             | Iva frutescens                         | 16   | 9                                      | 3  |
|                                   |                                | Distichlis spicata/<br>Spartina patens | 9  | 16                                     | 5  |
|                                   |                                | Scirpus Olneyi                         | 23   | 12                                     | 4  |
|                                   |                                | Polygonum hydropiperoides              | 3  | 6                                      | 2  |

TABLE 3 (Continued)

| <u>Vegetation Type</u>            | <u>n Sample Number</u> | <u>Species Present</u> | <u>Mean Cover (% 2-m<sup>2</sup> plot)</u> | <u>S.D. Standard Deviation</u> | <u>S.E. Standard Error of the Mean</u> |
|-----------------------------------|------------------------|------------------------|--|--------------------------------|--|
| Iva frutescens<br>(Western plots) | 12                     | Spartina cynosuroides  | 13   | 14                             | 4                                      |
|                                   |                        | Spartina alterniflora  | 3  | 5                              | 1                                      |
|                                   |                        | Solidago sempervirens  | 0.2  | 0.6                            | 0.2                                    |
|                                   |                        | Hibiscus palustris     | 0.8  | 3                              | 0.8                                    |
|                                   |                        | Pluchea camphorata     | 2  | 4                              | 1                                      |
|                                   |                        | Kosteletzkya virginica | 1  | 2                              | 0.7                                    |
|                                   |                        | Baccharis halimifolia  | 1  | 3                              | 0.8                                    |
|                                   |                        | Juncus spp.            | 0.8  | 2                              | 0.6                                    |
|                                   |                        | Panicum virgatum       | 0.8  | 3                              | 0.8                                    |
|                                   |                        | Mud/water              | 27   | 15                             | 4                                      |

# Mean Percent Cover: Iva Stands







FIGURE 30

## Mean Percent Cover

### Typha Stand



### Spartina / Distichlis Stand




FIGURE 31

# Percent Cover: Iva Plots







FIGURE 32

# Percent Cover: Iva Plots







FIGURE 33

# Percent Cover: Iva Plots

09







FIGURE 34

# Percent Cover: Iva Plots









FIGURE 35

# Percent Cover: Iva Plots







FIGURE 36

# Percent Cover: Iva Plots







FIGURE 37

# Percent Cover: Iva Plots

64






FIGURE 38

# Percent Cover: Typha Plots









FIGURE 39

# Percent Cover: Typha Plots







FIGURE 40

# Percent Cover: *Spartina patens* Plots









FIGURE 41

# Percent Cover: *Spartina patens* Plots












FIGURE 42

# Percent Cover: *Spartina patens* Plots





CORRELATIONS OF COVER, SPECTRAL SIGNATURE, AND PRODUCTIVITY  
IN HOG ISLAND MARSH

Successful correlations between marsh productivity and color infrared aerial photos have been made by Reimold (1971) and the results used for land management decisions. Reimold worked with a very large marsh of basically one community type: Spartina alterniflora. He found that the brighter and more saturated red colors on color infrared aerial transparencies were associated with more productive Spartina alterniflora areas in the marsh.

Since the marshes of Rhode River are small and heterogeneous, an attempt was made to use Reimold's method as an indicator of productivity differences between, rather than within plant communities. In Table 4, standing crop measurements taken in July and August 1971, for eight communities in Hog Island Marsh are compared with the color of these communities in Hog Island and Kirkpatrick Marshes and with the percent cover of the type species. The aerial transparencies used for this comparison were taken August 24, 1971 from an altitude of 3500 feet.

The correlation between standing crop and color saturation is ambiguous because of the characteristics of the different communities examined. In August, Typha angustifolia has dark brown flowering heads, Panicum virgatum has purplish brown flowering heads, and Phragmites communis has very dense, tan or purplish flowering heads. These flowering heads appear brown or brown-green on color infrared photos and consequently obscure bright green (red on color infrared) foliage underneath. There is understandably little correlation between cover and standing crop since small plants such as Spartina patens may form a dense cover while tall plants such as Typha angustifolia, with greater biomass per unit area, may form a sparse cover.

TABLE 4

| <u>Community</u>                                      | <u>Standing Crop</u><br><u>(g dry wt/m<sup>2</sup>)</u> <sup>1.</sup> | <u>Color on Color</u><br><u>Hog Island</u><br><u>Marsh</u> | <u>IR Photos</u><br><u>Kirkpatrick</u><br><u>Marsh</u> | <u>Mean Cover</u><br><u>(% 2-m<sup>2</sup> plot)</u> |
|---|---|--|--|--|
| <i>Phragmites communis</i>                            | 1114 ± 990  | bright pink  | bright red   |  |
| <i>Spartina cynosuroides</i>                          | 672 ± 300   | brown/red  | bright red   |  |
| <i>Typha angustifolia</i>                             | 626 ± 163   | brown/red  | brown/red  | 56   |
| <i>Spartina alterniflora</i>                          | 587 ± 158   | bright/dull<br>red   | bright/dull  |  |
| <i>Iva frutescens</i>                                 | 534 ± 177   | dull red   | dull red   | 25   |
| <i>Scirpus Olneyi</i>                                 | 472 ± 139   | tan red  | tan red  |  |
| <i>Spartina patens</i> /<br><i>Distichlis spicata</i> | 445 ± 125   | light pink   | light pink   | 70   |
| <i>Panicum virgatum</i>                               | 369 ± 104   | dull red/<br>brown   | dull red   |  |

1. From Drake and Hayes, unpublished data.

#### REFERENCES CITED

- Heller, R.C. , G.E. Doverspike, and R.C. Aldrich. 1964. Identification of tree species on large-scale panchromatic and color aerial photographs. U.S.D.A., Forest Service, Agriculture Handbook #261. 17 pp.
- Krumpe, P.F., H.R. DeSelm, and C.C. Amundsen. 1971. The delineation of forest cover and site parameters by multiband remote sensing. American Society of Photogrammetry, 37th annual meeting, March, 1971. Pp. 98-122.
- Reimold, R.J. 1971. Remote sensing of salt marsh productivity. Proc. Third Biennial Workshop on Color Aerial Photography in Plant Sciences. March 2-4, 1971. Gainesville, Fla. American Society of Photogrammetry. 13 pp.